

—20. *Parkinsonia aculeata*, L.—21. *Acacia*, sp. ?—22. *Piptadenia*? sp.—23. *Mimosa asperata*, L.—
24. *Turnera ulmifolia*, L.—25. *Sida rhombifolia*, L.—26. *Paritium tiliaceum*, St. Hil.

ALTERNANTHERA RADICATA, *Hook. fl.*; radice validâ lignosâ apice 3–5-cipiti, foliis stellatis recurvis lanuginosis teretibus carnosis elongato-linearibus integerrimis obtusis medio supernè canaliculatis, ramis floriferis radicalibus abbreviatis lanuginosis apice oligocephalis, capitulis globosis, perianthii foliolis oblongis obtusis emarginatis, antheris 2, stigmatibus bipartito.

Hab. Chatham Island, *C. Darwin, Esq.*

Radix pro plantâ maxima, elongata, lignosa, crassitie pollicis, supernè divisa. Folia perplurima, patienti-recurva, $1\frac{1}{2}$ –2 unc. longa, teretia, 1 lin. diametr., basi densè lanuginosa, ceterùm arachnoideo-tomentosa. Caules floriferi (seu scapi) foliis breviores, validi, supernè bracteati, simplices v. divisi, plerumque monocephali; bracteis 3–4, capitulo brevioribus, lanuginosis. Capitula alba, $\frac{1}{2}$ lin. lata, multiflora; floribus compressis lanâ subscariosâ immersis, $1\frac{1}{2}$ lin. longis. Bracteolæ floribus paulò longiores, laterales dorso acutæ, postica concava. Perianthii foliola enervia. Tubus stamineus cyathiformis, brevis; filamentis fertilibus 2, lateralibus, dilatatis, sterilibus nullis. Ovarium orbiculare, compressum. Stigmata brevia.

The root of this species is much larger than that of *A. subscaposa*, with which it agrees in habit. The foliage will distinguish it from that, and from every other described species.—*July 1847.*

ERRATA.

Page 194. No. 105 should be *PLANTAGO TOMENTOSA*, Lam.?

— 195. Before No. 114 dele *RUBIACEÆ*.

— 229. *CASTELA GALAPAGEIA*, *Hook. fl.* = varietas inermis *C. Nicholsonii*, Hook. The genus should be referred to *Simarubææ*.